

Determination of Aflatoxins in Medicinal Plants by High-Performance Liquid Chromatography–Tandem Mass Spectrometry

Nadeem A. Siddique^a, Mohd Mujeeb^a, Sayeed Ahmad^a, Bibhu P. Panda^b, Mohd Makhmoor^b

^a Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Jamia Hamdard (Hamdard University), New Delhi-110062, India. ^b Microbial and Pharmaceutical Biotechnology Laboratory, Faculty of Pharmacy, Jamia Hamdard (Hamdard University), New Delhi, India.

Received, June 1, 2013; Revised; Accepted, July 11, 2013; Published, July 14, 2013.

ABSTRACT – Purpose. The intention of the proposed work is to study the presence of the aflatoxins B₁, B₂, G₁ and G₂ in medicinal plants, namely *Mucuna pruriens*, *Delphinium denudatum* and *Portulaca oleraceae*. **Methodology.** The aflatoxins were extracted, purified by immunoaffinity column chromatography and analysed by high-performance liquid chromatography–tandem quadrupole mass spectrometry with electrospray ionisation (HPLC–MS/MS). Fungal count was carried out in PDA media. **Results.** A good linear relationship was found for AFB₁, AFB₂, AFG₁ and AFG₂ at 1–10 ppb ($r > 0.9995$). The analyte accuracy under three different spiking levels was 86.7–108.1 %, with low per cent relative standard deviations in each case. The aflatoxins can be separated within 5 to 7 min using an Agilent XDB C18-column. We found that AFB₁ and AFB₂ were in trace amounts below the detection limit in *M. pruriens* whilst they were not detected in *D. denudatum*. *P. oleraceae* was found to be contaminated with AFB₁ and AFB₂. AFG₁ and AFG₂ were not detected in *M. pruriens*, *P. oleraceae* and were below the detection limit in *D. denudatum*. This was consistent with very low numbers of fungal colonies observed after 6 hr of incubation. **Conclusion.** The analytical method developed is simple, precise, accurate, economical and can be effectively used to determine the aflatoxins in medicinal plants and therefore to control the quality of products. The aflatoxin levels in the plant extracts examined were related to the minimal fungal load in the medicinal plants examined.

This article is open to **POST-PUBLICATION REVIEW**. Registered readers (see “For Readers”) may **comment** by clicking on ABSTRACT on the issue’s contents page.

INTRODUCTION

The plants *Mucuna pruriens*, *Delphinium denudatum* and *Portulaca oleraceae* have the potential for wide therapeutic applications. *Mucuna pruriens* L (Velvet beans; Fabaceae), a herbaceous forage and food legume, has found widespread usage in the management of several diseases, including diabetes, rheumatoid arthritis, atherosclerosis, male infertility and nervous disorders, and is a good source of 3,4-dihydroxyphenylalanine (DOPA) (1, 2). In addition to health benefits the seeds possess potent aphrodisiac, geriatric tonic and vermifuge activities (3). The major constituent of *Mucuna* seeds was found to be L-dopa (5%), along with minor amounts of methylated and nonmethylated tetrahydroisoquinolines (0.25 %) (4). *Delphinium denudatum* Wall (Jadwar; Ranunculaceae) is one of the important medicinal drugs used as indigenous medicine in India, and the entire plant is reported to be useful in a variety of ailments (5). The root is used

in various medical formulations in Unani and Ayurveda to reduce the withdrawal symptoms in people on de-addiction therapy (6). It is a natural analgesic and is prescribed for relief from toothache (7). Herbalists recommend the roots in the treatment of fungal infections, asthma, cough, jaundice and nervous problems (8). It is considered as a natural blood purifier and prescribed for people who have digestive problems and provides relief from painful wounds and inflammation (9). Chemically *D. denudatum* contains many active compounds, some of which are alkaloids including delphocurarine, staphisagrine, delphine, condelphine, denudatin, talatizidine, acetylhetero-phylistine and a

Corresponding Author: Mohd Mujeeb, M.Pharm, Ph.D; Assistant Professor; Bio Active Natural Product Laboratory (Nano Building); Department of Pharmacognosy and Phytochemistry; Faculty of Pharmacy; Jamia Hamdard (Hamdard University); New Delhi, India; E-mail: mhdmjb@gmail.com

diterpenoid alkaloid identical to condelphine (10). *Portulaca oleracea* L. (Purslane; Portulacaceae) grows widely in different areas of the world including India. The seeds are regarded as valuable in the treatment of urinary problems, digestive problems and cardiovascular diseases (11). Studies demonstrated a variety of pharmacological activities, including hypoglycemic (12, 13), hypocholesterolemic, antioxidant, analgesic, anti-inflammatory, skeletal muscle relaxant, smooth muscle relaxant, neuroprotective (14), anxiolytic, sedative, anticonvulsant, antiulcerogenic, bronchodilatory, wound healing, and antifungal effects (15). It contains a variety of bioconstituents, including catecholamines, l-noradrenaline, dopamine, α -amyrin, β -amyrin and portuloside A (16, 17, 18).

Aflatoxins are a group of structurally related toxic metabolites of mycotoxins produced by *Aspergillus flavus* and *Aspergillus parasiticus* (19). In the aflatoxin group, about 16 compounds are known, but only aflatoxin B₁, B₂, G₁, G₂, M₁ and M₂ are routinely monitored (20, 21). Aflatoxin B₁, B₂, G₁ and G₂ are chemical derivatives of difurancoumarin (Fig. 1). Aflatoxin B₁ is the prevalent compound in plant samples and it is acutely toxic and the most carcinogenic amongst the aflatoxins (22). Aflatoxins may increase stress

susceptibility and compromise growth efficiency. The clinical signs of aflatoxicosis in humans are extremely varied. Signs of acute aflatoxicosis include depression, nervousness, abdominal pain, diarrhea and death (23). The aflatoxins occur widely in food, especially in corn, nuts, peanuts, coconut, fruits, dried fruits and beer (24, 25). High moisture and temperature are two main factors that cause the occurrence of mycotoxins at pre-harvest and post harvest stages (26). Fungi are widely distributed as environmental contaminants (27), in fact, under favorable conditions of temperature and humidity, they grow on many commodities including cereals, oil seeds, nuts, herb-teas and spices (28). They have also been found in medicinal herbs that are not fully dried or stored improperly (e.g. at high temperature and humidity) (29, 30). Despite many studies on mycotoxins in agricultural products, only a few are concerned with spices and herbal medicines. The determination of aflatoxins in herbs is becoming important for medicinal safety, as more and more people are using medicinal herbs for complementary and alternative therapy and they play an important role in the economy, especially in developing countries. Previously aflatoxins were analyzed in peanut, cocoa, pistachio, chili, ginger (31, 32), chocolate, cotton-seed, job's tears, pepper, paprika, feed and wheat samples (33).




Figure 1. Structure of (a) AFB₁ (b) AFB₂ (c) AFG₁ (d) AFG₂

The current acceptance level of total aflatoxin in maize set by the United States- Food and Drug Administration (USFDA) is 20 µg/kg (34). Health Canada has set acceptance criteria of < 20 µg/kg (ppb) for aflatoxins (B₁+B₂+G₁+G₂), and < 5 µg/kg (ppb) for aflatoxin B₁ for evening primrose oil, sugar cane, sugar beets, cottonseed, peanuts, corn, ginseng or any substance derived from these sources (35). Recently, the Codex Alimentarius Commission, the Joint FAO/WHO Food Standards Program, adopted a limit of 15µg/kg for total aflatoxins (36). In 2008, the Codex Alimentarius set a maximum level of 10 µg/kg total aflatoxins in ready-to-eat almonds, hazelnuts, and pistachios (37). In the European Union, the total aflatoxins and AFB₁ levels in human commodities are regulated with maximum residue levels (MRLs) that cannot be greater than 2 and 4 µg/kg, respectively (38). In the year 2002 the European Food Safety Authority (EFSA) set maximum limits in spices (*Capsicum* spp., *Piper* spp., *Myristicafragrans*, *Zingiber officinale*, *Curcuma longa*) (AFB₁ 5 µg/kg; total AFs 10 µg/kg) (02/472/EC) (39).

Environmentally sound and most widespread methods applicable for quantitative determination of aflatoxin content in different samples are thin-layer chromatography (40, 41) and high performance liquid chromatographic (HPLC) (42, 43). Aflatoxins can be separated and detected using either normal- or reversed-phase HPLC methods mainly with fluorescence spectrometric detection. The use of two-dimensional thin-layer chromatography (2D-TLC) (44, 45) and high performance thin-layer chromatography (HPTLC) for the determination of aflatoxins has been reported (46). An alternative complex media based method is reported to detect the natural fluorescence of aflatoxins released by the growing mycelium (47) or relies on multiplex polymerase chain reaction (PCR) and real time polymerase chain reaction (RT-PCR) detection of genes or their transcripts involved in the aflatoxin biosynthetic pathway (48, 49). An enzyme-linked immunosorbent assay (ELISA) for AFB₁, and levels of total aflatoxins were quantified and confirmed by liquid chromatography combined with tandem mass spectrometry (LC-MS/MS) (50). However, a lot of time and a tedious pre- or post-column derivatization is required with conventional HPLC in order to provide better detection limits of AFB₁, AFB₂, AFG₁ and AFG₂ (51). Furthermore, many other structurally similar components to the AFs in the plants can also

be retained by the aflatoxin immunoaffinity column, and as a result produce false positive results in many chromatographic techniques.

Despite the numerous uses and the various biologically active chemical constituents reported in *M. pruriens*, *D. denudatum* and *P. oleracea*, no data on aflatoxin detection of these plants are available in the literature. In order to ensure safety and quality we have collectively developed a cost efficient, flexible and rapid method for the quantification of aflatoxins in *M. pruriens*, *D. denudatum* and *P. oleracea* as these plants are commonly consumed by people without any awareness of their safety. There exist no well-accepted scientific methods for proper determination and quality control of aflatoxins in selected plant materials. The aim of this study was to screen the content of aflatoxins as well as providing a sensitive, accurate and reproducible analytical method for the detection of aflatoxins. Further, the purpose of this study is to provide useful information on aflatoxin levels in *M. pruriens*, *D. denudatum* and *P. oleracea* and to achieve a confirmatory method for quick determination of AFB₁, AFB₂, AFG₁ and AFG₂ by high performance liquid chromatography–electrospray ionization tandem mass spectrometry (HPLC–MS/MS). The proposed method overcame the drawbacks of HPLC by using tandem MS/MS detection, which can determine the presence of a specific substance more accurately than HPLC and avoids the tedious derivatization process. The method is validated for linearity, recovery, sensitivity, precision and repeatability, and was shown to successfully determine the levels of AFs in plants samples.

METHODS

Equipment and chemicals

The analysis was performed on an Agilent 1200 system using an Agilent XDB-C18 4.6-50 mm column containing 1.8 mm particles. MS/MS detection was achieved on an Agilent 6410B tandem quadrupole mass spectrometer using an electrospray ion (ESI) source in positive mode. Data were acquired and processed using the Masshunter software (Agilent). Acetonitrile, methanol, ammonium acetate and HPLC grade water were supplied by Merck (Darmstadt, Germany). AFB₁, AFB₂, AFG₁ and AFG₂ were purchased from Sigma (St. Louis, MO). All other inorganic chemicals and organic solvents were of reagent grade.

HPLC–MS/MS analysis

Chromatographic analyses were carried out using a binary gradient elution with 70:30 v/v methanol:acetonitrile as eluent A and 5 mM aqueous ammonium acetate as eluent B. The total run time was 5 to 8 min. The injection volume was 5 mL, the flow rate was set at 5 μ L/min and the column temperature was maintained at 30°C. The ionisation source of the MS/MS detector had 4.0 kV capillary voltage, 350 °C source temperature and 9 L/min desolvation gas flow (both gases were nitrogen).

Sample collection

All sample species were collected from a local market of New Delhi, India and were authenticated by Dr. H.B. Singh (Chief Scientist & Head, Raw materials Herbarium & Museum, NISCAIR, New Delhi). A voucher specimen (RHMD/1704/04) has been deposited in the Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Jamia Hamdard, New Delhi. All samples were ground to powders and kept in moisture proof paper bags to keep their water content constant before analysis. The herbs and their parts used are summarized in Table 1.

Sample preparation

The samples (5 g) were sonicated in 100 mL 70:30 v/v methanol:water for 45 min and centrifuged (Remi Compufuge-CR30) for 10 min, then 5 mL of the supernatant was drawn, diluted with 30 mL milli-Q water, and passed through the immunoaffinity column at no more than 5 mL/min (the column was previously conditioned with 30 mL distilled water). The column was rinsed with 15 mL double distilled water to remove the matrix components and then dried by passing air through to remove any adherent water. The eluate was diluted to 2 mL with water and passed through a 0.22 mm filter, and the filtrate was injected into the HPLC–MS/MS.

Peak recognition

The AFs in samples were recognized by comparing their retention times and MS spectra (MRM mode) with the standards. The standards used are summarized in Table 1.

Fungal count in the plant materials

All the plant materials were diluted with sterile water and 500 μ L of the samples were spread over potato dextrose agar (PDA)-Ox bile media. They were incubated for 5 days at 28 °C and 75 % relative humidity, and the colony forming units (CFU) of the fungal strains were observed.

RESULTS

Selection of mobile phase

The HPLC mobile phase was optimized by considering the retention time, peak response, sensitivity, peak sharpness as well broadening and separation efficiencies. Methanol–water in the combination of 60:40 v/v completely separated the analytes but was found not suitable for the column, while acetonitrile–water 60:30 v/v failed to separate the AFs. Methanol:acetonitrile in a combination of 70:30 v/v with 5.0 mM aqueous ammonium acetate was found the best mobile phase for AF standards separation (Fig. 2).

Optimisation of the mass spectra

Mass spectrometry detection was carried out on positive ionisation mode because this mode gave sharp and sensitive signals. Mass spectrometry detection was optimised by injecting the AF standards (100 ng/mL) under positive ionisation mode. The ion fragments were evaluated by their run time to the most abundant m/z and the ion with the uppermost intensity was selected as the basic ion for quantitation (Fig. 3).

Table 1. Summary of aflatoxins and herbs used in this study

Standard aflatoxin	Tested herb		
	Part used	Species	Family
B ₁			
B ₂	Seed	<i>Mucuna pruriens</i>	Fabaceae
G ₁	Root	<i>Delphinium denudatum</i>	Ranunculaceae
G ₂	Seed	<i>Portulaca oleraceae</i>	Portulacaceae




Figure 2. HPLC-MS/MS Chromatograms of standard mycotoxins comprising a mixture of aflatoxins (B₁, B₂, G₁ and G₂). Mobile phase; methanol: acetonitrile (70:30 v/v) eluent A and aqueous ammonium acetate (5 mM) eluent B. Total run time (5 to 8 min), injection volume (5 mL) and flow rate (5 μ L/min).




Figure 3. Mass spectral analysis of standard aflatoxins. (A) Std. AFB₁ run time (7.8-8.1) and m/z (313.0) [M+H]⁺ (B) AFB₂ run time (7.1-7.5) and m/z (315.0) [M+H]⁺ (C) AFG₁ run time (6.5-6.8) and m/z (329.0) [M+H]⁺ (D) AFG₂ run time (5.6-6.0) and m/z (331.0).

METHOD VALIDATION

Limits of detection and limits of quantification

The AF analyses showed good results in the MRM mode, the LOD reaching 0.28–1.10 μ g/kg and LOQ 0.79–3.34 μ g/kg, with a signal-to-noise ratio (S:N) of 3. Linear calibration curve for various concentrations of AFs ranging from 1 ppb–10 ppb (three points) were analysed. The calibration curves were constructed by plotting the response against the concentration. A linear relationship was obtained for each compound, and the correlation coefficients ranged from 0.9992 to 0.9995 (Table 2). The relative standard deviations (RSDs) of precision of AF mixtures were 7.04–2.17 % (n=6). Excellent retention times for AFB₁

(7.973 \pm 0.018), AFB₂ (7.361 \pm 0.017), AFG₁ (6.689 \pm 0.015) and AFG₂ (5.830 \pm 0.026) were observed for corresponding mixtures of AFs (Table 2).

Accuracy

Accuracy analysis was performed by testing samples at three different concentrations 1 ppb, 5 ppb and 10 ppb. All plants were used for the accuracy test to ensure that the method has wide adaptability. The plant sample (5 g) was spiked with high, medium and low levels of AF standards. The accuracy ranged from 86.7% to 108.1% (Table 3), indicating good accuracy of the method.

Determination of aflatoxins in plant samples

HPLC-MS/MS Chromatograms (Counts vs Acquisition time (min)) for AFB₁, AFB₂, AFG₁ and AFG₂ were determined for *M. pruriens* (Fig. 4), *D. denudatum* (Fig. 5) and *P. oleraceae* (Fig. 6). The study revealed mass peaks of AFB₁ and AFB₂ in *M. pruriens* below the detection limit whilst a resultant mass peak of AFB₁ and AFB₂ was observed in *P. oleraceae*, whereas no mass peak was found in *D. denudatum* (Figs. 4, 5 and 6). The aflatoxin levels in the plants investigated is summarized in Table 4.

Fungal count

Total fungal colonies observed per gram of *M. pruriens*, *D. denudatum* and *P. oleraceae* were 18×10^2 , 4×10^2 and 20×10^2 respectively in PDA-ox bile media. The total fungal CFU also show that there was very low growth of fungus in these tested plant materials, which is consistent with the low levels of mycotoxins.

DISCUSSION

An LC-MS method was developed which was found to be suitable for the rapid and accurate assessment of aflatoxin levels in selected plant samples. The method was subsequently applied to determine the natural occurrence of AFs in selected widely used commercial medicinal plant samples (Table 1). We found that *P. oleraceae* was contaminated with AFB₁ (1.675 µg/kg) and AFB₂ (1.335 µg/kg), whilst levels in *M. pruriens* were below the detection limit for AFB₁ (0.04 µg/kg) and AFB₂ (0.05 µg/kg) and were not detected in *D. denudatum*. AFG₁ and AFG₂ were absent in *M. pruriens* and *P. oleraceae*, whilst a relatively low value (0.07 µg/kg) of AFG₂ was found in *D. denudatum* (Table 4).

Table 2. Linearity, limit of detection, limit of quantification and precision for the analysis of AFB₁, AFB₂, AFG₁ and AFG₂ by HPLC-MS/MS. Data expressed as mean±SD. ^an=6; ^{b, b1}S/N=3 (Signal-to-noise ratio); ^cn=6. Abbreviations: RT: Retention Time; LOD: Limit of detection; LOQ: Limit of quantification; RSD: Relative standard deviations.

Compound	RT±SD	Regression Slope	Correlation Intercept	(LOD) _b Coefficient ^a	(LOQ) _b ¹ µg/kg	Precision ^c µg/kg	(RSD %)
				R ²			
B ₁	7.973±0.018	-44.8238	927.2519	0.9992	0.2849	0.7996	0.704988
B ₂	7.361±0.017	-11.4898	742.1691	0.9995	0.6903	2.0918	0.155408
G ₁	6.689±0.015	-180.806	787.3258	0.9859	0.0839	0.2545	0.048938
G ₂	5.830±0.026	-12.9685	277.9480	0.9991	1.1044	3.3468	0.217735

Table 3. Recovery of aflatoxins AFB₁, AFB₂, AFG₁ and AFG₂ in low level, medium level and high level, Data expressed as mean±SD. Abbreviations: L: Low level (1 ppb); M: Medium level (5 ppb); H: High level: (10 ppb), ^an=6.

Compound	Concentration (ppb)	Response ^a ±SD	Accuracy (%)
AFB ₁	L	955.66±5.13	108.1
AFB ₂	L	224.33±4.50	105.8
AFG ₁	L	665±4.0	107.4
AFG ₂	L	77±2.64	108.7
AFB ₁	M	4456.66±5.508	97.1
AFB ₂	M	1074.66±3.512	97.9
AFG ₁	M	3232.33±4.163	86.7
AFG ₂	M	391.66±3.055	96.9
AFB ₁	H	9286.66±5.132	100.6
AFB ₂	H	2224.66±4.509	100.5
AFG ₁	H	8158.33±5.033	105.9
AFG ₂	H	826.66±4.509	100.7




Figure 4. HPLC-MS/MS Chromatogram of *M. pururiens* Counts vs Acquisition time (min); (A) Resultant mass peak of AFB₁ observed below the detection limit (0.04 µg/kg), (B) Resultant mass peak of AFB₂ observed below the detection limit (0.05 µg/kg) (C) No resultant mass peak of AFG₁ observed (D) No resultant mass peak of AFG₂ observed. Mobile phase; methanol: acetonitrile (70:30 v/v) eluent A and aqueous ammonium acetate (5 mM) eluent B. Total run time (5 to 8 min), injection volume (5 mL) and flow rate (5 µL/min). The black line in the chromatogram is for quantifier ion and the blue line shows the qualifier ion for the MRM transition for LC-MS/MS of aflatoxins. Ratio in the chromatogram is quantifier to qualifier ion ratio for the compound (which determines confirmation of a compound).




Figure 5. HPLC-MS/MS Chromatogram of *D. denudatum*. Counts vs Acquisition time (min); (A) No resultant mass peak of AFB₁ observed (B) No resultant mass peak of AFB₂ observed (C) No Resultant mass peak of AFG₁ observed (D) Resultant mass peak of AFG₂ observed below the detection limit (0.07 µg/kg). Mobile phase: methanol: acetonitrile (70:30 v/v) eluent A and aqueous ammonium acetate (5 mM) eluent B. Total run time (5 to 8 min), injection volume (5 mL) and flow rate (5 µL/min). The black line in the chromatogram is for quantifier ion and blue line shows the qualifier ion for the MRM transition for LC-MS/MS of aflatoxins. Ratio in the chromatogram is quantifier to qualifier ion ratio for the compound, (which determines confirmation of a compound).




Figure 6. HPLC-MS/MS Chromatogram of *P. oleraceae* Counts vs Acquisition time (min) (A) Resultant mass peak of AFB₁ observed (1.675 µg/kg) (B) Resultant mass peak of AFB₂ observed (1.335 µg/kg) (C) No resultant mass peak of AFG₁ observed (D) No resultant mass peak of AFG₂ observed. Mobile phase: methanol: acetonitrile (70:30 v/v) eluent A and aqueous ammonium acetate (5 mM) eluent B. Total run time (5 to 8 min), injection volume (5 mL) and flow rate (5 µL/min). The black line in the chromatogram is for quantifier ion and the blue line shows the qualifier ion for the MRM transition for LC-MS/MS of aflatoxins. Ratio in the chromatogram is quantifier to qualifier ion ratio for the compound, (which determines confirmation of a compound).

Table 4. Summary of contamination levels of aflatoxins in herbs ($\mu\text{g}/\text{kg}$) obtained from LC-Mass spectroscopy
Abbreviations: BDL: Below detection limit; ND: Not detected.

Name of herb	AFB ₁	AFB ₂	AFG ₁	AFG ₂	Total AF
<i>Mucuna pruriens</i>	BDL (0.04)	BDL (0.05)	ND	ND	0.09
<i>Delphinium denudatum</i>	ND	ND	ND	BDL (0.07)	0.07
<i>Portulaca oleraceae</i>	1.675	1.335	ND	ND	3.01

Amongst the four AFs, AFB₁ was detected as the most common contaminant. The AFs occurred in the order of AFB₁>AFB₂>AFG₁>AFG₂. *Aspergillus flavus* is a very common fungus which infects plants and produces AFB₁ and AFB₂. However levels of AFG₁ and AFG₂ are less often detected in plants because the fungus *Aspergillus parasiticus* does not infect plants very often (52). This may account for the high incidence and concentrations of AFB₁ and AFB₂ compared with AFG₁ and AFG₂ in the three plants investigated in this study. Rhizomes, roots, seeds and fruits have apparently higher AFs than in flowers, indicating that the herbs rich in starch and oil might be more favorable for fungal growth (53). In particular, contamination levels of some samples i.e. root of *D. denudatum*, seeds of *M. pruriens* and *P. oleraceae* were notable, as all of these have a content of fat or fatty oil, suggesting that plants and their products rich in fat or fatty acids should be carefully monitored. The plant samples should be dried as soon as possible and stored in moisture-proof containers with good aeration to avert the growth of AFs, as inappropriate storage and moisture content in samples encourages augmentation of mycotoxins (54). We conclude from this study that the risk of AF contamination in medicinal herb like *M. pruriens*, *D. denudatum* and *P. oleraceae* appears low but the presence of AFs in these plants needs to be analyzed before processing to confirm the absence of mycotoxins. The consumer must be cautious while consuming these plants for medicinal purposes. The monitoring scheme described could be applied to determine the aflatoxin contamination in a large number of samples in a cost and time effective manner. The data obtained from this monitoring can be used as a basis for risk analysis of aflatoxin contamination, thereby maintaining the aflatoxin contents at the lowest possible levels.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the Department of Pharmacognosy and Phytochemistry, Faculty of

Pharmacy, Jamia Hamdard, New Delhi and Council of scientific and industrial research (CSIR) Government of India for financial assistance (09/591(0106)/2011-EMR-I). The authors thank Professor Basil Roufogalis for help with preparation of the manuscript.

REFERENCES

- Bhaskar A, Vidhya V, Ramya M. Hypoglycemic effect of *Mucuna pruriens* seed extract on normal and streptozotocin-diabetic rats. *Fitoterapia* 2008 Dec; 79(7-8):539-543.
- Siddhuraju P, Becker K. Rapid reversed-phase high performance liquid chromatographic method for the quantification of L-dopa (L-3,4-dihydroxyphenylalanine), nonmethylated and methylated tetrahydroisoquinoline compounds from *Mucuna* beans. *Food Chem* 2001 Feb; 72(3):389-394.
- Carsky RJ, Tarawali SA, Becker M, Chikoye D, Tian G, Sanginga N, et al. *Mucuna*- Herbaceous cover legume with potential for multiple uses. *Resource & Crop Management Research Monograph* 1998; 25:54.
- Spencer JPE, Jenner P, Halliwell B. Superoxide-dependent GSH depletion by L-DOPA and dopamine. Relevance to Parkinson's disease. *Neuro Report* 1995 Jul; 6(11):1480-1484.
- Qudsia N, Jafri MA. Unani drug, Jadwar (*Delphinium denudatum* Wall.) - A review. *IJTK* 2006 Oct; 5(4):463-467.
- Zafar S, Aftab MA, Siddiqui TA. Jadwar (*Delphinium denudatum* Wall.) Roots - A Boon in Unani Medicine. *Hamdard Med* 2003; 6(2):9-14.
- Rahman SZ, Khan RA, Kumar A. Experimental study of the morphine de-addiction properties of *Delphinium denudatum* Wall. *BMC Complem Altern Med* 2002 May; 2(6).
- Rahman SZ, Khan RA, Kumar A. Pharmacological study of *Delphinium denudatum* Wall in morphine de-addiction. *J Neurochem* 2007 Aug; 102 (Suppl.1):142-143.
- Ahmad G, Yousuf A, Khan KM, Tajuddin NA. The anti stress activity of a gem containing Unani formulation against diverse stress. *J Ethnopharmacol* 2004 Feb; 90(2-3):367-74.
- Singh N, Chopra KL. Diterpine alkaloid isolation and study of two new alkaloids. *J Pharma and Pharmaceutic* 1962; 14:288-293.

11. Oh KB, Chang IM, Hwang KJ, Mar W. Detection of antifungal activity in *Portulaca oleracea* by a single-cell bioassay system. *Phytother Res* 2000 Dec; 14(5):329-332.
12. Islam MW, Zakaria MNM, Radhakrishnan R, Habibullah M, Chan K. Evaluation of analgesic activity of the aerial parts of *Portulaca oleracea* v. *sativa* and its comparison with other two related species. *J Pharm Pharmacol* 1998; 50(Suppl):226.
13. Chan K, Islam MW, Kamil M, Radhakrishnan R, Zakaria MN, Habibullah M, Attas A, et al. The analgesic and anti-inflammatory effects of *Portulaca oleracea* L. subsp. *Sativa* (Haw.) Celak. *J Ethnopharmacol* 2000 Dec; 73(3):445-51.
14. Radhakrishnan R, Zakaria MN, Islam MW, Chen HB, Kamil M, Chan K, Al-Attas A, et al. Neuropharmacological actions of *Portulaca oleracea* L v. *sativa* (Haw). *J Ethnopharmacol* 2001 Jul; 76(2):171-176.
15. Ullah M O , Haque M , Urmi K F , Zulfiker A H , Anita E S , Begum M , Hamid K. Anti-bacterial activity and brine shrimp lethality bioassay of methanolic extracts of fourteen different edible vegetables from Bangladesh. *Asian Pac J Trop Biomed* 2013 Jan; 3(1):1-7.
16. Li HY, Wang XS. A study on extracting method of polysaccharide from *Portulaca oleracea*. *J Branch Campus First Military Med* 2005; 2:186-187.
17. Mohammad TB, Mohammad HB, Farhad M. Antitussive effect of *Portulaca oleracea* L in guinea pigs. *Iran J Pharmaceut Res* 2004;3: 187-190.
18. Meng FB, Wu RG. Appraisal on medicinal values of *Portulaca oleracea* L. *Forest Investig Des* 2008; 1:77-78.
19. Eaton DL.; Groopman JD.; The Toxicology of aflatoxins, Human health, veterinary and agricultural significance. Academic Press, New York, 1994.
20. Hussein HS, Brasel JM. Toxicity, metabolism and impact of mycotoxins on humans and animals. *J Toxicol* 2001; 167:101-109.
21. Joshua H. Determination of aflatoxins by reversed-phase high-performance liquid chromatography with post-column in-line photochemical derivatization and fluorescence detection. *J Chromatogr A* 1993 Nov; 654 (2):247-254.
22. Abeywickrama E, Bean GA. Toxicogenic *Aspergillus flavus* and aflatoxins in Sri Lankan medicinal plant material. *Mycopathologia* 1991 Mar; 113(3):187-190.
23. Ip SP, Che CT. Determination of aflatoxins in Chinese medicinal herbs by high-performance liquid chromatography using immunoaffinity column cleanup Improvement of recovery. *J Chromatogr A* 2006 Dec; 1135 (2):241-244.
24. Escobar A, Regueiro OS. Determination of aflatoxin B1 in food and feedstuffs in Cuba using an immunoenzymatic reagent kit (Aflacen). *J Food Prot* 2002; 65:219-221.
25. Mphande FA, Siame BA, Taylor JE. Fungi, aflatoxins and cyclopiazonic acid associated with peanut retailing in Botswana. *J Food Prot* 2004 Jan; 67(1):96-102.
26. Urraca JL, Marazuela MC, Moreno-Bondi MC. Analysis for zearalenone and a-zearalenol in cereals and swine feed using accelerated solvent extraction and liquid chromatography with fluorescence detection. *Anal Chim Acta* 2004 Oct; 524 (1-2):175-183.
27. Juan C, Zinedine A, Molto JC, Idrissi L, Manes J. Aflatoxins levels in dried fruits and nuts from Rabat-Sale area, Morocco. *Food Control* 2008 Sept; 19 (9):849-853.
28. Gilbert J, Anklam E. Validation of analytical methods for determining mycotoxins in foodstuffs. *Trends Anal Chem* 2002 Jun; 21(6-7): 468-486.
29. Kerstin H, Charity M. Aflatoxin control and prevention strategies in key crops of Sub-Saharan Africa. *Afr J Microbiol Res* 2011; 5(5):459-466.
30. Reddy KRN, Salleh B. Co-Occurrence of Moulds and Mycotoxins in Maize Grains Used for Animal Feeds in Malaysia. *J Anim Vet Adv* 2011 Mar; 10(5):668-673.
31. Roy AK, Sinha KK, Chourasia HK. Aflatoxin contamination of some common drug plants. *Appl Environ Microbiol* 1988 Mar; 54(3):842-843.
32. Han Z, Zheng Y, Luan L, Cai Z, Ren Y, Wu Y, et al. An ultra-high-performance liquid chromatography-tandem mass spectrometry method for simultaneous determination of aflatoxins B1, B2, G1, G2, M1 and M2 in traditional Chinese medicines. *Anal Chim Acta* 2010 Apr; 664(2):165-171.
33. Sugita-Konishi Y, Sato T, Saito S, Nakajima M, Tabata S, Tanaka T, Norizuki H, Itoh Y, Kai S, Sugiyama K, Kamata Y, Yoshiike N, Kumagai S et al. Exposure to aflatoxins in Japan: risk assessment for aflatoxin B1. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 2010; 27(3):365-72.
34. Ubwa ST, Asemave K, Igbum GO. Preliminary screening of aflatoxin level in maizen (*Zea mays* L.) in some selected markets in Benue State, Nigeria. *AJPAC* 2012 Dec; 6(14):159-163.
35. The Natural Health Products Directorate (NHPD), Quality of Natural Health Products Guide". Version 3.0- 17 August, 2012: http://www.hc-sc.gc.ca/dhp-mps/consultation /natur/consult_quality-qualit eng.php.
36. Codex Alimentarius Commission. Joint FAO/WHO food standards programme, codex committee on food additives and contaminants. Thirty-third session (2001) CODEX, Hague, Netherlands.
37. Safety evaluation of certain food additives and contaminants. WHO Food Additive Series: 59 (2008). <http://www.who.int/foodsafety/chem/jecfa/publications/monographs/en/index.html>.
38. European Economic Community Council. Commission regulation (EC) No.1525/98. Official Journal of European Communities 1998; 201:43-46.

39. Commission Regulation (EC) No 472/2002 of 12 March 2002 amending regulation (EC) No 466/2001 setting maximum levels for certain contaminants in foodstuffs. Official Journal of the European Communities L75/18, Luxembourg.
40. Alexander RF, Baur MC. Note on a two TLC procedure for determining Aflatoxin in maize. Cereal Chem 1977 May; 54:699-704.
41. Younis YMH, Malik MK. TLC and HPLC assay of aflatoxin contamination in Sudanese peanuts and peanut products. Kuwait J Sci Eng 2003; 30 (1):79-93.
42. Waltking AE, Wilson D. Liquid chromatographic analysis of aflatoxin using post-column photochemical derivatization: collaborative study. J AOAC Int 2006 May; 89(3):678-692.
43. Sobolev VS. Simple, rapid, and inexpensive cleanup method for quantitation of aflatoxins in important agricultural products by HPLC. J Agric Food Chem 2007 Mar; 55(6):2136-2141.
44. Kamkar A. A study on the occurrence of aflatoxin M1 in raw milk produced in Sarab city of Iran. Food Control 2005 Sept; 16(7):593-599.
45. Alborzi S, Pourabbas B, Rashidi M, Astaneh B. Aflatoxin M1 contamination in pasteurized milk in Shiraz (south of Iran). Food Control 2006 Jul; 17(7):582-584.
46. Heshmati A, Milani JM. Contamination of UHT milk by aflatoxin M1 in Iran. Food Control 2010 Jan; 21(1):19-22.
47. Maragos CM, Appell M, Lippolis V, Visconti A, Catucci L, Pascale M. Use of cyclodextrins as modifiers of fluorescence in the detection of mycotoxins. Food Addit Contam 2008; 25(2):164-171.
48. Somashekar D, Rati ER, Chandrashekar A. PCR-restriction fragment length analysis of *aflR* gene for differentiation and detection of *Aspergillus flavus* and *Aspergillus parasiticus* in maize. Int J Food Microbiol 2004 May; 93(1):101-107.
49. Scherm B, Palomba M, Serra D, Marcello A, Migheli Q. Detection of transcripts of the aflatoxin genes *aflD*, *aflO*, and *aflP* by reverse transcription-polymerase chain reaction allows differentiation of aflatoxin-producing and non-producing isolates of *Aspergillus flavus* and *Aspergillus parasiticus*. Int J Food Microbiol 2005 Feb; 98(2):201-210.
50. Shim WB, Kim K, Ofori JA, Chung YC, Chung DH. Occurrence of aflatoxins in herbal medicine distributed in South Korea. J Food Protect 2012; 75(11):1991-99.
51. Ali N, Hashim NH, Saad B, Safan K, Nakajima M, Yoshizawa T, et al. Evaluation of a method to determine the natural occurrence of aflatoxins in commercial traditional herbal medicines from Malaysia and Indonesia. Food Chem Toxicol 2005 Dec; 43(43):1763-1772.
52. Dorner JW, Richard JC, Diener UL. The relationship of *Aspergillus flavus* and *A. parasiticus* with reference to production of aflatoxins and cyclopiazonic acid. Mycopathologia 1984 Aug; 87(1-2):13-15.
53. Romagnoli B, Menna V, Gruppioni N, Bergamini C. Aflatoxins in spices, aromatic herbs, herb-teas and medicinal plants marketed in Italy Food control 2007 June; 18(6):697-701.
54. Schroeder HW. Factors influencing the development of aflatoxins in some field crops. J Stored Prod Res 1969 Nov; 5(3):187-192.